

Starvation Induced Changes in Quantitative Measures of Liver Function in the Rat

B. A. Hansen* and H. E. Poulsen†

*Department of Medicine A, Division of Hepatology and Laboratories of Experimental Pathology, Rigshospitalet A 2152, DK-2100 Copenhagen, Denmark.

†Department of Pharmacology, University of Copenhagen, DK-2100 Copenhagen, Denmark.
(Reprint requests to B.A.H.)

ABSTRACT Liver function was estimated quantitatively as the capacity of urea-N synthesis (CUNS), the galactose elimination capacity (GEC) and the antipyrine plasma clearance (APC) in rats following 24, 48, and 96 h of food deprivation. CUNS rose to a maximum following 24 h of starvation, then decreased to the control value at 48, 72, and 96 h ($p = 0.02$). GEC decreased following starvation, significant so at 48 h ($p = 0.02$). APC decreased at 48, and 72 h ($p = 0.01$), but rose to the control value at 96 h. It is concluded that a precise standardisation of starvation is important for interpretation of minor changes in CUNS, GEC, and APC.

INTRODUCTION

The capacity of urea-N synthesis (CUNS), the galactose elimination capacity (GEC), and the antipyrine plasma clearance (APC) are all established measures of 'the functional liver mass' [1].

Urea cycle enzymes have increased activity during states when protein catabolism is increased, such as starvation [2]. Hepatic antipyrine metabolism is found to be influenced by malnutrition [3], while the effect of starvation on galactose elimination to our knowledge has only been investigated following 18 h of food deprivation [4].

According to this the liver function tests could be changed following various degrees of food deprivation, meaning that a precise standardisation with regard to nutritional status should be carefully taken into consideration. The aim of the present study was to investigate the effect of starvation for various periods on liver function tests in the rat.

MATERIAL AND METHODS

Experimental design

Measurement of the capacity of urea-N synthesis (CUNS), the galactose elimination capacity (GEC), and the antipyrine plasma clearance (APC) was made simultaneously in 25 female Wistar rats.

The animals were fed rat pellets and kept at constant ambient temperature with a fixed 12 h controlled light/dark cycle. For determination of CUNS, GEC, and APC anesthesia was induced by intraperitoneal injection

of thiopental, 100 mg per kg body weight. Estimation of CUNS, GEC and APC was made in fed rats and following 24, 48, 72, and 96 h of starvation. Five animals were examined in each group. Polyethylene catheters were inserted into both jugular veins for infusion, and into the right common carotid artery for blood sampling, immediately after tracheotomy and intubation. The kidney vessels were ligated after exteriorisation via the dorsal route. Alanine was administered as a single injection of 0.40–0.75 ml of a 1.09 mol/l solution in sterile water, followed by a constant infusion for 70 min at 1.5–3.0 ml/h of 224 mmol/l solution by means of a roller pump (Perfusor Secura). Steady-state amino acid concentration was defined as less than 10% change during a period of 50 min or longer. Simultaneously galactose (Kabi, Sweden) was given as an intravenous priming dose (200 μ mol/100 g body weight) followed by a constant infusion (1.0–1.5 μ mol/min).

Immediately before administration of alanine and galactose blood samples were drawn into chilled tubes containing aprotinin and EDTA and stored at -20°C until analysis of insulin and glucagon concentration by RIA. Glucagon was extracted by ethanol precipitation according to Heding [5]. Samples (150 μ l) were taken

Table 1 Body weight (BW)g, and liver weight (LW)g at various intervals of starvation. Values are mean \pm SEM of five animals

	Food deprivation for				
	Fed	24 h	48 h	72 h	96 h
BW	222 \pm 8	211 \pm 4	216 \pm 7	207 \pm 5	187 \pm 3
LW	7.7 \pm 0.2	6.6 \pm 0.2	6.4 \pm 0.2	6.1 \pm 0.1	5.7 \pm 0.3

Table 2 The capacity of urea-N synthesis (CUNS) $\mu\text{mol}(\text{min}\cdot 100\text{ g BW})^{-1}$, galactose elimination capacity (GEC) $\mu\text{mol}(\text{min}\cdot 100\text{ g BW})^{-1}$, and the antipyrine plasma clearance (APC) $\text{ml}(\text{min}\cdot 100\text{ g BW})^{-1}$ at various intervals of starvation. Values are mean \pm SEM of five animals

	Food deprivation for				
	Fed	24 h	48 h	72 h	96 h
CUNS	8.2 \pm 1.0	11.2 \pm 0.6	9.1 \pm 0.5	6.8 \pm 1.2	7.3 \pm 0.7
GEC	1.26 \pm 0.08	1.07 \pm 0.20	0.81 \pm 0.15	1.04 \pm 0.23	0.95 \pm 0.25
APC	0.34 \pm 0.03	0.27 \pm 0.02	0.22 \pm 0.01	0.23 \pm 0.02	0.33 \pm 0.06

Table 3 Plasma glucose (mmol/l), glucagon (ng/l), and insulin (mU/l) at various intervals of starvation. Values are mean \pm SDM of five animals

	Food deprivation for				
	Fed	24 h	48 h	72 h	96 h
Glucose	5.8 \pm 0.3	4.2 \pm 0.2	3.5 \pm 0.3	3.5 \pm 0.2	3.2 \pm 0.4
Glucagon	80 \pm 26	204 \pm 30	100 \pm 42	79 \pm 17	90 \pm 19
Insulin	45.9 \pm 5.5	19.5 \pm 5.6	21.2 \pm 5.0	19.1 \pm 6.1	22.3 \pm 1.8

after an equilibration period of 20 min, at intervals of 10 min, for determination of urea [6], total alpha-amino-N [7] and galactose concentrations [8]. For determination of APC, the animals received antipyrine 1 ml (4 mg/ml) by gastic tube 5 h prior to the procedures described above. At the time of investigation blood samples were taken and heparinised plasma stored at -20°C until analysis for antipyrine by HPLC [9].

Calculation. CUNS $\mu\text{mol}(\text{min}\cdot 100\text{ g BW})^{-1}$ was calculated as

$$\text{CUNS} = (dc_u/dt) \cdot 0.63 \text{ BW} \cdot 1.25$$

where dc_u/dt is the slope of the linear regression of arterial blood urea nitrogen concentration on time during steady state, 0.63 BW is the estimated volume of distribution of urea [10], and 1.25 is a correction for intestinal hydrolysis of urea [11]. GEC $\mu\text{mol}(\text{min}\cdot 100\text{ g BW})^{-1}$ was calculated as

$$\text{GEC} = I - (dc/dt \cdot 0.40 \text{ BW})$$

where I is the infusion rate of galactose, dc/dt is the linear slope of the galactose blood concentration-time curve, and 0.40 BW is the volume of distribution of galactose [4].

APC $\text{ml}(\text{min}\cdot 100\text{ g BW})^{-1}$ was calculated as

$$\text{APC} = (\ln(D/0.66 \text{ BW}) - (\ln(c_i))/t$$

where D is the antipyrine dose given, 0.66 BW is the antipyrine volume of distribution [9], and c_i is the concentration corresponding to the sampling time t [9]. The period of 5 h between antipyrine dose and the determination of the plasma concentration corresponds to the optimal time for sampling [9]. The liver was removed, blotted on filter paper, and weighed.

Statistical calculations. A two-sided t-test was performed between each of the treatment group versus the

control group, resulting in $p < 0.05$ as level of significance.

RESULTS

Table 1 shows the decrease in body weight and liver during 96 h of starvation ($p = 0.001$, $p = 0.001$, respectively).

Table 2 and Figure 1 show the sequential changes in CUNS, GEC, and APC at various intervals of starvation. CUNS rose to a maximum following 24 h of starvation, then decreased to the control value at 48, 72, and 96 h ($p = 0.02$). GEC decreased following starvation, significant so at 48 h ($p = 0.02$). APC decreased at 48 and 72 h ($p = 0.01$), but rose to the control value at 96 h.

Table 3 shows the sequential changes in plasma concentration of glucose, glucagon and insulin at various intervals of starvation. The glucose concentration decreased during starvation, and was nearly half the control value after 96 h ($p = 0.001$). The glucagon response to starvation was at maximum at 24 h ($p = 0.01$), then decreased to the control value after 48, 72, and 96 h. The insulin concentration was only half the control value at 24, 48, 72, and 96 h ($p = 0.01$).

DISCUSSION

The present study demonstrates that both the capacity of urea-N synthesis (CUNS), the galactose elimination capacity (GEC), and the antipyrine plasma clearance




Fig. 1 The capacity of urea-N synthesis (CUNS) (panel a), galactose elimination capacity (GEC) (panel b), and antipyrine plasma clearance (APC) (panel c) at various intervals of starvation. Symbols indicate mean, bars SEM of five animals.

(APC) is changed during starvation. As shown previously by others, also the plasma concentrations of glucose, glucagon, and insulin were changed [12].

The method used for estimation of CUNS presuppose substrate concentration independent conditions [11], meaning that starvation induced changes in the amino acid concentration can be neglected. The liver function tests used are given relative to body weight, as starvation induced changes in both body and liver weight.

One key event in the transition from the fed to the fasted state is an increase in gluconeogenesis [13]. Considering urea synthesis as an index of such an event, the

increase in CUNS found at 24 h of starvation, is in parallel with gluconeogenesis after this interval. We found that starvation for longer than 48 h decreased CUNS, in accordance with inhibited protein break down after prolonged starvation [14]. In vitro, Schimke found the urea cycle enzyme activity increased following starvation for 4 and 7 days [2]. As urea excretion in that situation was decreased, it apparently has no significance in vivo, or might be due to regulation by circulating humoral factors, not present in the in vitro system.

Recently it has been demonstrated that within short periods CUNS is regulated by insulin [15] whereas glucagon has a time dependent stimulating effect [16]. The increase in CUNS in the 24–48 h fasted rats can therefore be attributed to the decreased circulating plasma insulin and increased glucagon concentrations.

In accordance with the findings of Keiding [4], there was no change in GEC during starvation for 24 h. Following starvation for 48 h we found a decrease in GEC, meaning that the rate limiting enzyme in galactose metabolism, galactokinase [4], could be under influence of the metabolic alteration induced by food deprivation.

APC is decreased considerably in protein-caloric malnutrition and is reestablished after nutritional rehabilitation [3]. In the present study, APC was decreased following 48 and 72 h of starvation. At 96 h APC was not different from that of the control. This could be interpreted as an adaptation to starvation condition, but the mechanism is not clear.

It is concluded that it is important also in human studies to use a precise standardisation with regard to starvation during determination of urea synthesis rate, galactose elimination capacity, and antipyrine clearance, in order to make a correct interpretation of minor changes in these measurements.

ACKNOWLEDGEMENTS

Our thanks are due to technician Bjorg Krog for expert laboratory assistance. We are indebted to Liselotte Hansen and Kirsten Prisholm for performing the insulin analysis.

Novo (Denmark) generously supplied iodinated insulin and glucagon, and antisera. The Lundbæk foundation is thanked for donation of HPLC apparatus.

REFERENCES

- [1] Hansen B A, Poulsen H E 1986 The capacity of urea-N synthesis as a quantitative measure of the liver mass in rats. *Journal of Hepatology* 2: 468–474
- [2] Schimke R T 1962 Differential effects of fasting and protein-free diets on levels of urea cycle enzymes in rat. *The Journal of Biological Chemistry* 237: 1921–1924
- [3] Tranvouez J L, Lerebours E, Chretien P et al 1985 Hepatic antipyrine metabolism in malnourished patients: influence of the type of malnutrition and course after

- nutritional rehabilitation. *The American Journal of Clinical Nutrition* 41: 1257-1264
- [4] Keiding S 1973 Galactose elimination capacity in the rat. *Scandinavian Journal of Clinical and Laboratory Investigation* 31: 319-325
- [5] Heding L 1971 Radioimmunological determination of pancreatic and gut glucagon in plasma. *Diabetologia* 7: 10-19
- [6] Fawcett J K, Scott J E 1960 A rapid and precise method for determination of urea. *Journal of Clinical Pathology* 13: 156-159
- [7] Goodwin J F 1970 Spectrophotometric quantitation of plasma and urinary amino nitrogen with flurodinitrobenzene. *Standard Method Clinical Chemistry* 6: 89-98
- [8] Kurz G, Wallemfels K 1970 UV-test mit galactose-dehydrogenase, In: Bergmeyer, H U (eds). *Methoden der enzymatischen analyse*. Chemie, Weinheim/Bergstr
- [9] Pilsgaard H, Poulsen H E 1984 A one-sample method for antipyrine clearance determination in rats. *Pharmacology* 29: 110-116
- [10] Foy J, Schnieden H 1960 Estimation of total body water (virtual tritium space) in the rat, cat, rabbit, guinea-pig, and man, and of the biological half-life of tritium in man. *Journal of Physiology* 154: 169-176
- [11] Hansen B A, Vilstrup H 1985 A method for determination of the capacity of urea synthesis in the rat. *Scandinavian Journal of Clinical and Laboratory Investigation* 45: 315-320
- [12] Mlekusch W, Paletta B, Truppe W et al 1981 Plasma concentrations of glucose, corticosterone, glucagon and insulin and liver content of metabolic substrates and enzymes during starvation and additional hypoxia in the rat. *Hormone and Metabolic Research* 13: 612-6-4
- [13] Zorzano A, Herrera E 1983 Liver and kidney cortex gluconeogenesis from L-alanine in fed and starved rats. *International Journal of Biochemistry* 16: 263-267
- [14] Goodman M N, Ruderman N B 1980 Starvation in the rat. I. Effect of age and obesity on organ weights, RNA, DNA, and protein. *American Journal of Physiology* 238: E269-E276
- [15] Hansen B A, Krog B, Vilstrup H 1986 Insulin and glucose decreases the capacity of urea-N synthesis in the rat. *The Scandinavian Journal Clinical and Laboratory Investigation*. In press
- [16] Petersen K F, Hansen B A, Vilstrup H 1986 Time dependent stimulating effect of glucagon on the capacity of urea-N synthesis in rats. *Hormone and Metabolic Research*. In press